

In the specification:

◇◇ Please amend the title as follows:

Methods for inhibiting ~~modulating hematopoiesis and~~ vascular growth

◇◇ Please amend the first complete paragraph on page 1 as follows:

This application is a continuation of United States Serial No. 09/021,660 (U.S. Patent No. 6,713,065), filed February 10, 1998, which claims priority from provisional application Serial Ser. No. 60/037,513, filed ~~Feb.~~ February 10, 1997 and provisional application Serial Ser. No. 60/049,763, filed ~~Jun.~~ June 16, 1997, the disclosures of which are hereby incorporated by reference.

◇◇ Please amend the 2<sup>nd</sup> complete paragraph on page 6 as follows:

Figs. 1 A through 1D show ~~shows~~ four expression cassettes used to generate transgenic mice. (A) -179 lacZ $\epsilon\mu$ LCR (MB70) is the basic construct with the minimal  $\epsilon$ -globin promoter ( $\epsilon$ -pro), extending to -179 with respect to the start site of transcription (+1) and ~20 bp of  $\epsilon$ -globin 5'-untranslated region (shown as small black box below +1).  $\epsilon$ -pro is linked to a LacZ expression cassette containing a Kozak consensus sequence and translational start site (SDK region). Downstream from the LacZ reporter gene is a portion of the 3'-region of the  $\epsilon$ -globin gene containing part of exon 2, all of intron 2 (IVS 2) and all of exon 3; these sequences are shown as black boxes (exons) and a black line (intron). The 3'-untranslated region (containing the polyadenylation site, pA) is shown as a striped line. A truncated version of the LCR (the  $\mu$ LCR) is located downstream from the  $\epsilon$ -lacZ sequences. (B) -849 lacZ $\epsilon\mu$ LCR (MB73), (C)  $\epsilon$ -PRE(II+V)lacZ $\epsilon\mu$ LCR (MB72); and (D) -2kblacZ $\epsilon\mu$ LCR (MB92). A-D contain  $\epsilon$ -pro and different portions of the upstream regulatory region of the human  $\epsilon$ -globin gene. Eukaryotic sequences in A-D were excised from the vector by digestion with KpnI and NotI and then purified for microinjection into the male pronuclei of mouse zygotes.

◇◇ Please amend the 3<sup>rd</sup> complete paragraph on page 6 as follows:

Figs. 2 A through 2F show ~~shows~~ LacZ expression, correlated with the appearance of a blue stain, in primitive erythroblasts. (A) ~~(a)~~ is a diagrammatic representation of a 7.5 dpc embryo; (B) ~~(b)~~ is a transgenic embryo stained with XGal and viewed by bright field microscopy,

(Ce) is the same embryo viewed by dark field microscopy; ~~B-(a) (D-E) is a non-transgenic mouse; (b) and (e) are embryos at 8.5 dpc stained with XGal; C(a) (F) is a wild type 12.5 dpc embryo, (b) and (e) are transgenic 12.5 dpc embryos.~~ (1) wild type; (2) transgenic; (3) ectoplacental cone; (4) blood islands; (5) amniotic cavity; (6) trophectoderm; (7) allantois; (8) extraembryonic mesoderm; (9) embryo proper (epiblast).

◇◇ Please amend the 1<sup>st</sup> complete paragraph on page 7 as follows:

Figs. 3A through 3D show ~~shows~~ the formation of yolk sac-like structures by cultured blastocysts: (A) (a) transgenic blastocysts prior to culture; (B) (b) ~~Saesac~~-like structure (non transgenic) stained with benzidine to reveal hemoglobin containing cells; (C) (c) ~~Sae sac~~ from cultured transgenic blastocysts stained with XGal to reveal hemoglobin containing cells after 9 days of cultivation; and (D) (d) ~~Normal normal~~ 8.5 dpc transgenic embryo and yolk sac stained with XGal.

◇◇ Please amend the 5<sup>th</sup> complete paragraph on page 7 as follows:

Figs. 7-1A through 7-1D show ~~shows~~ that when transgenic explants of gastrulating embryos (isolated at 6.25-6.5dpc) are cultured on filters or glass slides for 72 hours, induction of embryonic hematopoiesis occurs in whole embryo, but is absent in epiblasts only, as determined by XGal staining. Dashed lines were drawn around the epiblasts to facilitate visualization of structures. ~~7A1 (7-1A)~~ whole embryo on a filter; ~~7A2 (7-1B)~~ epiblast on a filter; ~~7A3 (7-1C)~~ whole embryo on a slide; and ~~7A4 (7-1D)~~ epiblast on a slide.

◇◇ Please amend the 1<sup>st</sup> complete paragraph on page 8 as follows:

Figs. 7-2A through 7-2E show ~~7-1 shows~~ blood formation in transgenic embryonic explant cultures. (lacZ stained sections of embryos). Frozen tissue sections were XGAL stained to reveal cluster of lacZ-positive hematopoietic cells in the whole embryos (7-2A) 7B1, epiblasts (7-2B) 7B2, posterior embryo portions (7-2C, 7-2D), ~~7B3, 7B4~~ and transgenic anterior epiblast portion adjacent to the VE (7-2E), ~~7B5~~, but not in surrounding visceral endoderm and undifferentiated mesoderm nor in the nontransgenic VE tissue of anterior/VE recombinants. ~~(e)~~

◇◇ Please amend the 2<sup>nd</sup> complete paragraph on page 8 as follows:

Figs. 8-1A through 8-1C ~~show~~ shows induction of hematopoiesis by visceral endoderm (VE) signals. A (8-1A) dark-field photomicrograph of recombinant containing transgenic (Tg) epiblast and non-Tg VE showing localized lac Z staining in the embryo adjacent to the visceral endoderm; B (8-1B) schematic diagram corresponding to panel 1Aa. Abbreviations: Tg , transgenic; Ve, visceral endoderm; EryP, primitive erythroid cells; C (8-1C) bright field photomicrograph of recombinant shown in A 1A.

◇◇ Please amend the 3<sup>rd</sup> complete paragraph on page 8 as follows:

Fig. 8-2 8D shows induction of embryonic hematopoiesis in whole embryo, and in epiblast plus visceral endoderm, but none in epiblasts only, using RT-PCR. (All samples were prepared following a 72 hour *in vitro* incubation of embryos isolated at 6.5 dpc). Actin served as an internal control.

◇◇ Please amend the 1<sup>st</sup> complete paragraph on page 9 as follows:

Figs. 14A through 14F ~~show~~ shows the early development of the mouse at 2.5 dpc (A), 3.5 dpc (B), 4.5dpc (C), 5.5 dpc (D), 6.5 dpc (E), and 7.5 dpc (F). The region of early blood island formation occurs in the exocoelomic cavity (see F f) between the epiblast below which is surrounded by the visceral endoderm and the extraembryonic tissue above.

◇◇ Please amend the 2<sup>nd</sup> complete paragraph on page 9 as follows:

Figs. 15A through 15C ~~show~~ shows the experimental scheme for separation of epiblast into anterior and posterior portions. (A) depicts the entire 6.75 dpc embryo with visceral endoderm around the perimeter of the epiblast and the extraembryonic mesoderm. (B) depicts the embryo after the visceral endoderm has been stripped off and (C) shows the epiblast only, with a dotted line of transection showing how the anterior and posterior sections are physically divided before separate cultivation.

◇◇ Please amend the 3<sup>rd</sup> complete paragraph on page 9 as follows:

Figs. 16-1 shows that hematopoietic mesoderm arises from the posterior primitive streak (posterior mesoderm) when anterior and posterior portions of lacZ transgenic embryos are harvested at mid- to late-gastrulation. Top Panel-A: No staining is detected in anterior epiblasts.

Bottom Panel ~~B~~: Dark blue XGAL histochemical staining shows blood formation in cultured posterior epiblasts. Scale bar, 1mm.